

TREATMENT PACKS

HELPING TO GET YOU TO YOUR RIGHT DOSE

XCOPRI® (cenobamate tablets) CV is a prescription medicine used to treat partial-onset seizures in adults 18 years of age and older.

IMPORTANT SAFETY INFORMATION FOR XCOPRI® (cenobamate tablets) CV

DO NOT TAKE XCOPRI IF YOU:

- Are allergic to cenobamate or any of the other ingredients in XCOPRI.
- Have a genetic problem (called Familial Short QT syndrome) that affects the electrical system of the heart.

Please see additional Important Safety Information and full Prescribing Information and Medication Guide in pocket.



XCOPRI™
(cenobamate tablets) CV
12.5 • 25 • 50 • 100 • 150 • 200 mg

When you begin taking XCOPRI® (cenobamate tablets) CV, you will be prescribed a treatment pack at the lowest dose. Then, your dose should slowly increase every 2 weeks until you reach a maintenance dose that is right for you.

Please take XCOPRI exactly how your healthcare provider prescribes it. The recommended starting dosage for XCOPRI is 12.5 mg once daily, and the recommended maintenance dosage is 200 mg once daily. In some cases, your healthcare provider may increase this, if needed. However, your daily dose should not exceed 400 mg. If you have mild or moderate liver issues, your healthcare provider will make sure you do not exceed 200 mg daily. Use of XCOPRI in patients with severe liver issues is not recommended.

IMPORTANT SAFETY INFORMATION (cont'd)

Allergic reactions: XCOPRI can cause serious skin rash or other serious allergic reactions which may affect organs and other parts of your body like the liver or blood cells. You may or may not have a rash with these types of reactions.

Please see additional Important Safety Information and full Prescribing Information and Medication Guide in pocket.



Treatment packs help make it easier when your healthcare provider increases your dose

When your healthcare provider prescribes XCOPRI® (cenobamate tablets) CV, you will receive treatment packs of XCOPRI tablets to help you increase your dose. Your first pack contains 12.5 mg and 25 mg tablets, as shown here. Each treatment pack comes with detailed dosing instructions from your healthcare provider. It is important to follow these instructions carefully.



IMPORTANT SAFETY INFORMATION (cont'd)

Call your healthcare provider right away and go to the nearest emergency room if you have any of the following: swelling of your face, eyes, lips, or tongue, trouble swallowing or breathing, a skin rash, hives, fever, swollen glands, or sore throat that does not go away or comes and goes, painful sores in the mouth or around your eyes, yellowing of your skin or eyes, unusual bruising or bleeding, severe fatigue or weakness, severe muscle pain, frequent infections, or infections that do not go away. **Take XCOPRI exactly as your healthcare provider tells you to take it. It is very important to increase your dose of XCOPRI slowly, as instructed by your healthcare provider.**

Remember, do not stop taking XCOPRI® (cenobamate tablets) CV without first talking with your healthcare provider.

You can get treatment packs through **SK life science navigator Patient Services** with assistance from your healthcare provider.

Visit XCOPRI.com/SKLSINavigator.com to get started.

Fight against partial-onset seizures with XCOPRI.
Learn more at XCOPRI.com.

Please see additional Important Safety Information and full Prescribing Information and Medication Guide in pocket.

IMPORTANT SAFETY INFORMATION FOR XCOPRI® (cenobamate tablets) CV

DO NOT TAKE XCOPRI IF YOU:

- Are allergic to cenobamate or any of the other ingredients in XCOPRI.
- Have a genetic problem (called Familial Short QT syndrome) that affects the electrical system of the heart.

XCOPRI CAN CAUSE SERIOUS SIDE EFFECTS, INCLUDING:

Allergic reactions: XCOPRI can cause serious skin rash or other serious allergic reactions which may affect organs and other parts of your body like the liver or blood cells. You may or may not have a rash with these types of reactions. Call your healthcare provider right away and go to the nearest emergency room if you have any of the following: swelling of your face, eyes, lips, or tongue, trouble swallowing or breathing, a skin rash, hives, fever, swollen glands, or sore throat that does not go away or comes and goes, painful sores in the mouth or around your eyes, yellowing of your skin or eyes, unusual bruising or bleeding, severe fatigue or weakness, severe muscle pain, frequent infections, or infections that do not go away. **Take XCOPRI exactly as your healthcare provider tells you to take it. It is very important to increase your dose of XCOPRI slowly, as instructed by your healthcare provider.**

QT shortening: XCOPRI may cause problems with the electrical system of the heart (QT shortening). Call your healthcare provider if you have symptoms of QT shortening including fast heartbeat (heart palpitations) that last a long time or fainting.

PREGNANCY AND LACTATION: XCOPRI® (cenobamate tablets) CV may cause your birth control medicine to be less effective. **Talk to your health care provider about the best birth control method to use.**

Talk to your health care provider if you are pregnant or plan to become pregnant. It is not known if XCOPRI will harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking XCOPRI. You and your healthcare provider will decide if you should take XCOPRI while you are pregnant. If you become pregnant while taking XCOPRI, talk to your healthcare provider about registering with the North American Antiepileptic Drug (NAAED) Pregnancy Registry. The purpose of this registry is to collect information about the safety of antiepileptic medicine during pregnancy. You can enroll in this registry by calling 1-888-233-2334 or go to www.aedpregnancyregistry.org.

Talk to your health care provider if you are breastfeeding or plan to breastfeed. It is not known if XCOPRI passes into breastmilk. Talk to your healthcare provider about the best way to feed your baby while taking XCOPRI.

COMMON SIDE EFFECTS: The most common side effects in patients taking XCOPRI include dizziness, sleepiness, headache, double vision, and feeling tired.

These are not all the possible side effects of XCOPRI. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. For more information, ask your healthcare provider or pharmacist. **Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088** or at www.fda.gov/medwatch.

DRUG ABUSE: XCOPRI is a federally controlled substance (CV) because it can be abused or lead to dependence. Keep XCOPRI in a safe place to prevent misuse and abuse. Selling or giving away XCOPRI may harm others and is against the law.

INDICATION:

XCOPRI is a prescription medicine used to treat partial-onset seizures in adults 18 years of age and older.

It is not known if XCOPRI is safe and effective in children under 18 years of age.

Please see additional patient information in the Medication Guide. This information does not take the place of talking with your healthcare provider about your condition or your treatment.

Please see full Prescribing Information and Medication Guide in pocket.

QT shortening: XCOPRI® (cenobamate tablets) CV may cause problems with the electrical system of the heart (QT shortening). Call your healthcare provider if you have symptoms of QT shortening including fast heartbeat (heart palpitations) that last a long time or fainting.

Suicidal behavior and ideation: Antiepileptic drugs, including XCOPRI, may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. Call your health care provider right away if you have any of the following symptoms, especially if they are new, worse, or worry you: thoughts about suicide or dying; attempting to commit suicide; new or worse depression, anxiety, or irritability; feeling agitated or restless; panic attacks; trouble sleeping (insomnia); acting aggressive; being angry or violent; acting on dangerous impulses; an extreme increase in activity and talking (mania); or other unusual changes in behavior or mood.

Nervous system problems: XCOPRI may cause problems that affect your nervous system. Symptoms of nervous system problems include: dizziness, trouble walking or with coordination, feeling sleepy and tired, trouble concentrating, remembering, and thinking clearly, and vision problems. **Do not drive, operate heavy machinery, or do other dangerous activities until you know how XCOPRI affects you.**

Do not drink alcohol or take other medicines that can make you sleepy or dizzy while taking XCOPRI without first talking to your healthcare provider.

DISCONTINUATION: Do not stop taking XCOPRI without first talking to your healthcare provider. Stopping XCOPRI suddenly can cause serious problems. Stopping seizure medicine suddenly in a patient who has epilepsy can cause seizures that will not stop (status epilepticus).

DRUG INTERACTIONS: XCOPRI may affect the way other medicines work, and other medicines may affect how XCOPRI works. **Do not start or stop other medicines without talking to your healthcare provider.** Tell healthcare providers about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.



©2020 SK Life Science, Inc., a subsidiary of
SK Biopharmaceuticals Co., Ltd.
PM-US-XCOP-0010 03/20

